Yamini Maddala, M.D., Deepti Dhavaleshwar, M.D.,

Phone: 469.697.5100

Suprep Split Dose Bowel Preparation for Colonoscopy

Physician:

Purchase your Suprep Bowel Preparation Kit from the Pharmacy

Colonoscopy

Colonoscopy is a test that allows your doctor to look at the inner lining of your large intestine (rectum and colon). In order to achieve an accurate evaluation of your lower gastrointestinal (GI) tract, it is important that you properly prepare for the procedure. Doctors are able to obtain the clearest view of the bowel when it is flushed completely free of waste material. The thoroughness of this intestinal cleaning depends on you. Without your cooperation, the examination cannot accomplish its purpose and may have to be repeated.

Procedure Date:	_
Check In Time:	
Location:	

Digestive Health Center of Allen

981 State Highway 121, Suite 2100, Allen, TX 75013 214-383-8210

Methodist Craig Ranch Surgery Center

6045 Alma Rd Suite 100, McKinney, TX 75070 469-907-1650

Baylor of McKinney

5252, W. University Dr, Highway 380 at Lake Forest Drive, McKinney, TX 75071 469-764-1000

Medical Center of McKinney

4500 Medical Center Drive, McKinney, TX 75069 972-547-8000

Medical City Surgery Center McKinney

4510 Medical Center Drive, Suite 150, McKinney, TX 75069 972-547-1580

Texas Health Presbyterian Hospital Allen

1105 Central Expressway North, Allen, TX 75013 972-747-1000

Five (5) Days before the Procedure:

(See attached prescription, if a sample was not given).

- Do Not Drink Alcohol for (5) DAYS prior to your procedure.
- Do Not take any or ibuprofen products. Taking Tylenol (acetaminophen) is allowed
- Stop taking anticoagulants or blood thinners such as: Coumadin, (Warfarin), Celebrex, NSAIDS and Plavix. *If necessary we will request Cardiac clearance*

Day before Procedure:

- Follow a clear liquid diet throughout the day. Items include: water, clear
 juices (apply and white grape only) without pulp, clear soda, chicken or beef
 broth, popsicles, Jell-O, tea, coffee, etc. If you are diabetic, drink regular
 (non-diet) liquids in moderation throughout the day and only take half your
 dose of diabetes meds(pills or insulin)
- Avoid Food, dairy products, orange juice and anything that is RED or PURPLE in color.
 - **Step 1**: At **5 PM** Pour ONE (1) 6-ounce bottle of Suprep liquid into the cup.
 - **Step 2**: Add cool drinking water to the 16-ounce line on the cup and mix.
 - **Step 3**: Slowly drink ALL the liquid in the cup.
 - **Step 4**: *MOST IMPORTANT:* You MUST drink two (2) more 16-ounce cups of water over the next hour.
 - **Step 5:** At **9 PM:** Repeat steps 1 through 4 using the other 6-ounce bottle of Suprep
- You may continue to have clear liquids until bedtime.
- Be sure and finish all of the solution and water.
- Both 6-ounce bottles are required for a complete prep.



Step 1:

Pour **ONE** (1) 6-ounce bottle of SUPREP liquid into the mixing container



Step 2:

Adding cool drinking water to the 16-ounce line on the water container and mix.

Note: Be sure to dilute SUPREP as shown before you drink it.



Step 3:

Drink ALL the liquid in the container



Step 3:

You must drink two (2) 16-ounce containers of water over the next 1 hour

Note: You must finish drinking the final glass of water at least 2 hours, or as directed before your procedure.

For directions visit: www.yaminimaddalamd.com/patientinformation/directions-to-endoscopy/

Yamini Maddala, M.D., Deepti Dhavaleshwar, M.D.,

Phone: 469.697.5100

Day Of Procedure:

- At Repeat steps 1 through 4 using the other 6 ounce bottle of Suprep bowel preparation kit along with the 16-ounce container.
- DO NOT eat or drink anything 4 hours prior to your procedure. Except for any currently prescribed heart or blood pressure medications with a sip of water.
- If you are diabetic: To avoid dangerously low blood sugar, Do Not take your diabetic medication (pills or insulin) the morning of your procedure. You can resume your diabetes medications after the procedure once you have eaten.
- Plan on being at the facility for approximately 2-3 hours and ensure to arrive at the facility at least 1 hour before the scheduled procedure.
- A Driver must accompany you at the facility; they must stay in the facility the entire time of your procedure.

YOU MUST HAVE SOMEONE DRIVE YOU HOME
- No Taxi, Uber, Lyft or Bus!

IF A PRESCRIPTION IS SENT IN PLEASE PICK IT UP WITHIN 7 DAYS

Please call the office within 48 hours to Cancel or Reschedule your procedure to avoid a \$100 charge. If you have any questions please call Leslie at 469-697-5100